arXiv:1403.7759v1 [math.CV] 30 Mar 2014

Evaluations of some terminating hypergeometric ${}_2F_1(2)$ series

Y. S. Kim[a], A. K. Rathie[b] and R. B. Paris[c]

[a] Department of Mathematics Education, Wonkwang University, Iksan, Korea
E-Mail: yspkim@wonkwang.ac.kr
[b] Department of Mathematics, Central University of Kerala, Kasaragad 671123, Kerala, India
E-Mail: akrathie@cukerala.edu.in
[c] School of Engineering, Computing and Applied Mathematics, University of Abertay Dundee, Dundee DD1 1HG, UK
E-Mail: r.paris@abertay.ac.uk

Abstract

Explicit expressions for the hypergeometric series ${}_2F_1(-n,a;2a\pm j;2)$ and ${}_2F_1(-n,a;-2n\pm j;2)$ for positive integer n and arbitrary integer j are obtained with the help of generalizations of Kummer's second and third summation theorems obtained earlier by Rakha and Rathie. Results for $|j|\leq 5$ derived previously using different methods are also obtained as special cases.

Mathematics Subject Classification: 33C15, 33C20

Keywords: Terminating hypergeometric series, generalized Kummer's second and third summation theorems

1. Introduction

In a problem arising in a model of a biological problem, Samoletov [9] obtained by means of a mathematical induction argument the following sum containing factorials

$$\sum_{k=0}^{n}\frac{(-1)^k}{(n-k)!}\,\frac{(2k+1)!!}{k!(k+1)!}=\frac{(-1)^n}{\sqrt{n!(n+1)!}}\left(\sqrt{n+1}\;\frac{(n-1)!!}{n!!}\right)^{(-1)^n},$$

where throughout n denotes a positive integer and, as usual,

$$(2n)!!=2\cdot 4\cdot 6\cdots=2^n n!,\qquad (2n+1)!!=1\cdot 3\cdot 5\cdots=\frac{(2n+1)!}{2^n n!}.$$

Samoletov also expressed the above sum in the equivalent hypergeometric form

$${}_2F_1\left[\begin{matrix}-n,\frac{3}{2}\\2\end{matrix};2\right]=\begin{cases}\dfrac{\Gamma(\frac{1}{2}n+\frac{1}{2})}{\sqrt{\pi}\Gamma(\frac{1}{2}n+1)} & (n\text{ even})\\[2ex] \dfrac{-\Gamma(\frac{1}{2}n+1)}{\sqrt{\pi}\Gamma(\frac{1}{2}n+\frac{3}{2})} & (n\text{ odd})\end{cases}$$

Subsequently, Srivastava [12] pointed out that this result could be easily derived from a known hypergeometric summation formula [7, Vol. 2, p. 493] for ${}_2F_1(-n,a;2a-1;2)$ with $a=\frac{3}{2}$, which is a contiguous result to the well-known summation

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ 2a \end{array};2\right] = \frac{2^n\sqrt{\pi}\Gamma(1-a)}{(2a)_n\Gamma(\frac{1}{2}-\frac{1}{2}n)\Gamma(1-a-\frac{1}{2}n)} \qquad (n=0,1,2,\ldots). $$

The aim in this note is to obtain explicit expressions for

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ 2a\pm j \end{array};2\right] \qquad \text{and} \qquad {}_2F_1\left[\begin{array}{c} -n, a \\ -2n\pm j \end{array};2\right] \tag{1.1} $$

for arbitrary integer j. We shall employ the following generalizations of Kummer's second and third summation theorems given in [8] (we correct a misprint in Theorem 6 of this reference). These are respectively

$$ {}_2F_1\left[\begin{array}{c} \alpha,\beta \\ \frac{1}{2}(\alpha+\beta\pm j+1) \end{array};\frac{1}{2}\right] = \frac{\sqrt{\pi}\Gamma(\frac{1}{2}\alpha+\frac{1}{2}\beta+\frac{1}{2}\pm\frac{1}{2}j)}{\Gamma(\frac{1}{2}\alpha+\frac{1}{2})\Gamma(\frac{1}{2}\beta+\frac{1}{2})}\,\frac{\Gamma(\frac{1}{2}\alpha-\frac{1}{2}\beta+\frac{1}{2}\mp\frac{1}{2}j)}{\Gamma(\frac{1}{2}\alpha-\frac{1}{2}\beta+\frac{1}{2}+\frac{1}{2}j)} $$

$$ \times\sum_{r=0}^{j}(\mp1)^r\binom{j}{r}\,\frac{(\frac{1}{2}\beta)_{r/2}}{(\frac{1}{2}\alpha+\frac{1}{2})_{(r-j)/2}} \tag{1.2} $$

and

$$ {}_2F_1\left[\begin{array}{c} \alpha, 1-\alpha\pm j \\ \gamma \end{array};\frac{1}{2}\right] = \frac{2^{\pm j}\Gamma(\frac{1}{2}\gamma)\Gamma(\frac{1}{2}\gamma+\frac{1}{2})}{\Gamma(\frac{1}{2}\gamma+\frac{1}{2}\alpha)\Gamma(\frac{1}{2}\gamma-\frac{1}{2}\alpha+\frac{1}{2})}\,\frac{\Gamma(\alpha\mp j)}{\Gamma(\alpha+\epsilon_j)} $$

$$ \times\sum_{r=0}^{j}(\mp1)^r\binom{j}{r}\,\frac{(\frac{1}{2}\gamma-\frac{1}{2}\alpha)_{r/2}}{(\frac{1}{2}\gamma+\frac{1}{2}\alpha)_{r/2-\delta_j}} \tag{1.3} $$

for $j=0,1,2,\ldots$, where $\epsilon_j=0$ (resp. j), $\delta_j=j$ (resp. 0) when the upper (resp. lower) signs are taken and $(a)_k=\Gamma(a+k)/\Gamma(a)$ is the Pochhammer symbol defined for *arbitrary* index k. When $j=0$, the summations (1.2) and (1.3) reduce to the well-known second and third summation theorems due to Kummer [11, p. 243]

$$ {}_2F_1\left[\begin{array}{c} \alpha,\beta \\ \frac{1}{2}(\alpha+\beta+1) \end{array};\frac{1}{2}\right] = \frac{\sqrt{\pi}\Gamma(\frac{1}{2}\alpha+\frac{1}{2}\beta+\frac{1}{2})}{\Gamma(\frac{1}{2}\alpha+\frac{1}{2})\Gamma(\frac{1}{2}\beta+\frac{1}{2})} $$

and[1]

$$ {}_2F_1\left[\begin{array}{c} \alpha, 1-\alpha \\ \gamma \end{array};\frac{1}{2}\right] = \frac{\Gamma(\frac{1}{2}\gamma)\Gamma(\frac{1}{2}\gamma+\frac{1}{2})}{\Gamma(\frac{1}{2}\gamma+\frac{1}{2}\alpha)\Gamma(\frac{1}{2}\gamma-\frac{1}{2}\alpha+\frac{1}{2})}. $$

In addition, we shall make use of the transformation [6, (15.8.6)]

$$ {}_2F_1\left[\begin{array}{c} -n,\beta \\ \gamma \end{array};2\right] = \frac{(-2)^n(\beta)_n}{(\gamma)_n}\,{}_2F_1\left[\begin{array}{c} -n, 1-\gamma-n \\ 1-\beta-n \end{array};\frac{1}{2}\right]. \tag{1.4} $$

Expressions for the series in (1.1) for arbitrary integer j have recently been obtained by Chu [2] using a different approach. This involved expressing the series for $j\neq0$ as finite sums of ${}_2F_1(2)$ series in (1.1) with $j=0$. The cases with $|j|\leq5$ have also been given by Kim and Rathie [3] and Kim *et al.* [4]. An application of the first series in (1.1) for $j=0,1,\ldots,5$ has been discussed in [5].

[1] In [11, p. 243], this summation formula is referred to as Bailey's theorem. However, it has been pointed out in [1] that this theorem was originally found by Kummer.

2. Statement of the results

Our principal results are stated in the following two theorems.

Theorem 1 *Let n be a positive integer, a be a complex parameter and define $j_0 = \lfloor \frac{1}{2}j \rfloor$. Then we have*

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a \pm j \end{array}; 2\right] = \frac{2^{2n}(\frac{1}{2})_n}{(2a \pm j)_{2n}} \sum_{r=0}^{j_0} (-1)^r \binom{j}{2r} (-n)_r (a+\delta_j)_{n-r} \tag{2.1} $$

and

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a \pm j \end{array}; 2\right] = \frac{\pm 2^{2n}(\frac{3}{2})_n}{(2a \pm j)_{2n+1}} \sum_{r=0}^{j_0} (-1)^r \binom{j}{2r+1} (-n)_r (a+\delta_j)_{n-r} \tag{2.2} $$

for $j = 0, 1, 2, \ldots$, where $\delta_j = j$ (resp. 0) when the upper (resp. lower) signs are taken.

Proof. From the result (1.4) we have

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ 2a \pm j \end{array}; 2\right] = \frac{(-2)^n (a)_n}{(2a \pm j)_n} \, {}_2F_1\left[\begin{array}{c} -n, 1-2a \mp j - n \\ 1-a-n \end{array}; \frac{1}{2}\right]. $$

The hypergeometric function on the right-hand side can be summed by the generalized second Kummer summation theorem (1.2), where we put $\alpha = 1 - 2a - j - n$ and $\beta = -n$. After some straightforward algebra we obtain

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ 2a + j \end{array}; 2\right] = \frac{2^n \sqrt{\pi}}{(2a+j)_n \Gamma(-\frac{1}{2}n)\Gamma(-\frac{1}{2}n + \frac{1}{2})} \sum_{r=0}^{j} (-1)^r \binom{j}{r} \frac{\Gamma(-\frac{1}{2}n + \frac{1}{2}r)}{(1-a-j)_{(r-n)/2}} $$

and

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ 2a - j \end{array}; 2\right] = \frac{2^n \sqrt{\pi}}{(2a-j)_n \Gamma(-\frac{1}{2}n)\Gamma(-\frac{1}{2}n + \frac{1}{2})} \sum_{r=0}^{j} \binom{j}{r} \frac{\Gamma(-\frac{1}{2}n + \frac{1}{2}r)}{(1-a)_{(r-n)/2}}. $$

Changing n to $2n$ and $2n+1$ and using the properties of the gamma function and

$$ (a)_{2n} = 2^{2n}(a)_n (a + \tfrac{1}{2})_n, \qquad (a)_{2n+1} = 2^{2n} a (\tfrac{1}{2}a + \tfrac{1}{2})_n (\tfrac{1}{2}a + 1)_n, $$

we then find the results stated in the theorem. □

We remark that in (2.2) the upper limit of summation can be replaced by $\lfloor \frac{1}{2}j \rfloor - 1$ when j is even. Also, since $(-n)_r$ vanishes when $r > n$, it is possible to replace the upper summation limit in both (2.1) and (2.2) by n whenever $n > \lfloor \frac{1}{2}j \rfloor$.

Theorem 2 *Let n be a positive integer and a be a complex parameter. Then we have*

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n + j \end{array}; 2\right] = \frac{2^{2n-j}(n-j)!}{(2n-j)!} \sum_{r=0}^{j} \binom{j}{r} (\tfrac{1}{2}a + \tfrac{1}{2} - \tfrac{1}{2}r)_n \tag{2.3} $$

provided j does not lie in the interval $[n+1, 2n]$ where the hypergeometric function on the left-hand side of (2.3) is, in general, not defined, and

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n - j \end{array}; 2\right] = \frac{2^{2n+j} n!}{(2n+j)!} \sum_{r=0}^{j} (-1)^r \binom{j}{r} (\tfrac{1}{2}a + \tfrac{1}{2} - \tfrac{1}{2}r)_{n+j} \tag{2.4} $$

for $j = 0, 1, 2, \ldots$. When $j \geq 2n+1$ in (2.3), the ratio of factorials $(n-j)!/(2n-j)!$ can be replaced by $(-1)^n (j-2n-1)!/(j-n-1)!$.

Proof. From the result (1.4) we have

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n \mp j \end{array};2\right] = \frac{2^n (a)_n (n \pm j)!}{(2n \pm j)_n}\, {}_2F_1\left[\begin{array}{c} -n, 1+n \pm j \\ 1-a-n \end{array};\frac{1}{2}\right] $$

when the parameters are such that the hypergeometric functions make sense. The hypergeometric function on the right-hand side can be summed by the generalized third Kummer theorem (1.3), where we put $\alpha = -n$ and $\gamma = 1-a-n$. Some straightforward algebra using the properties of the gamma function then yields the results (2.3) and (2.4) in the theorem. □

The sums on the right-hand sides of (2.3) and (2.4) can be written in an alternative form involving just two Pochhammer symbols containing the index n by making use of the result

$$ (\alpha - r)_n = \frac{(\alpha)_n (1-\alpha)_r}{(1-\alpha-n)_r} $$

for positive integers r and n. Then we find, with $j_0 = \lfloor \frac{1}{2} j \rfloor$,

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n+j \end{array};2\right] $$

$$ = \frac{2^{2n-j}(n-j)!}{(2n-j)!}\left\{ (\tfrac{1}{2}a+\tfrac{1}{2})_n \sum_{r=0}^{j_0} \binom{j}{2r} A_r(n,0) + (\tfrac{1}{2}a)_n \sum_{r=0}^{j_0} \binom{j}{2r+1} B_r(n,0) \right\} \quad (2.5) $$

and

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n-j \end{array};2\right] $$

$$ = \frac{2^{2n+j} n!}{(2n+j)!}\left\{ (\tfrac{1}{2}a+\tfrac{1}{2})_{n+j} \sum_{r=0}^{j_0} \binom{j}{2r} A_r(n,j) - (\tfrac{1}{2}a)_{n+j} \sum_{r=0}^{j_0} \binom{j}{2r+1} B_r(n,j) \right\}, \quad (2.6) $$

where

$$ A_r(n,j) := \frac{(\frac{1}{2} - \frac{1}{2}a)_r}{(\frac{1}{2} - \frac{1}{2}a - n - j)_r}, \qquad B_r(n,j) := \frac{(1 - \frac{1}{2}a)_r}{(1 - \frac{1}{2}a - n - j)_r}. $$

Again, when j is even, the upper summation limit in the second sums in (2.5) and (2.6) can be replaced by $j_0 - 1$, if so desired.

3. Special cases

If we set $0 \le j \le 5$ in (2.1) we obtain the following summations:

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{1}{2})_n} = {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a+1 \end{array};2\right], \quad (3.1) $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a+2 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{3}{2})_n}\left(1 + \frac{2n}{a+1}\right), \quad (3.2) $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a+3 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{3}{2})_n}\left(1 + \frac{4n}{a+2}\right), \quad (3.3) $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a+4 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{5}{2})_n}\left(1 + \frac{8n}{a+2} + \frac{16n(n-1)}{(a+2)(a+3)}\right), \quad (3.4) $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a+5 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a+\frac{5}{2})_n}\left(1+\frac{12n}{a+3}+\frac{16n(n-1)}{(a+3)(a+4)}\right) \tag{3.5} $$

and

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a-1 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a-\frac{1}{2})_n}, \tag{3.6} $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a-2 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a-\frac{1}{2})_n}\left(1+\frac{2n}{a-1}\right), \tag{3.7} $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a-3 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a-\frac{3}{2})_n}\left(1+\frac{4n}{a-1}\right), \tag{3.8} $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a-4 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a-\frac{3}{2})_n}\left(1+\frac{8n}{a-2}+\frac{8n(n-1)}{(a-1)(a-2)}\right), \tag{3.9} $$

$$ {}_2F_1\left[\begin{array}{c} -2n, a \\ 2a-5 \end{array};2\right] = \frac{(\frac{1}{2})_n}{(a-\frac{5}{2})_n}\left(1+\frac{12n}{a-2}+\frac{16n(n-1)}{(a-1)(a-2)}\right). \tag{3.10} $$

Similarly, if we set $0 \leq j \leq 5$ in (2.2) we obtain the following summations:

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a \end{array};2\right] = 0, \tag{3.11} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a+1 \end{array};2\right] = \frac{(\frac{3}{2})_n}{(2a+1)(a+\frac{3}{2})_n}, \tag{3.12} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a+2 \end{array};2\right] = \frac{2(\frac{3}{2})_n}{(2a+2)(a+\frac{3}{2})_n}, \tag{3.13} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a+3 \end{array};2\right] = \frac{(\frac{3}{2})_n}{(2a+3)(a+\frac{5}{2})_n}\left(3+\frac{4n}{a+2}\right), \tag{3.14} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a+4 \end{array};2\right] = \frac{(\frac{3}{2})_n}{(2a+4)(a+\frac{5}{2})_n}\left(4+\frac{8n}{a+3}\right), \tag{3.15} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a+5 \end{array};2\right] = \frac{(\frac{3}{2})_n}{(2a+5)(a+\frac{7}{2})_n}\left(5+\frac{20n}{a+3}+\frac{16n(n-1)}{(a+3)(a+4)}\right) \tag{3.16} $$

and

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a-1 \end{array};2\right] = -\frac{(\frac{3}{2})_n}{(2a-1)(a+\frac{1}{2})_n}, \tag{3.17} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a-2 \end{array};2\right] = -\frac{2(\frac{3}{2})_n}{(2a-2)(a-\frac{1}{2})_n}, \tag{3.18} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a-3 \end{array};2\right] = -\frac{(\frac{3}{2})_n}{(2a-3)(a-\frac{1}{2})_n}\left(3+\frac{4n}{a-1}\right), \tag{3.19} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a-4 \end{array};2\right] = -\frac{(\frac{3}{2})_n}{(2a-4)(a-\frac{3}{2})_n}\left(4+\frac{8n}{a-1}\right), \tag{3.20} $$

$$ {}_2F_1\left[\begin{array}{c} -2n-1, a \\ 2a-5 \end{array};2\right] = -\frac{(\frac{3}{2})_n}{(2a-5)(a-\frac{3}{2})_n}\left(5+\frac{20n}{a-2}+\frac{16n(n-1)}{(a-1)(a-2)}\right). \tag{3.21} $$

Finally, from (2.5) and (2.6) we obtain:

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n \end{array};2\right] = \frac{2^{2n}n!}{(2n)!}(\tfrac{1}{2}a+\tfrac{1}{2})_n = \frac{(\frac{1}{2}a+\frac{1}{2})_n}{(\frac{1}{2})_n}, \tag{3.22} $$

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n+1 \end{array};2\right] = \frac{2^{2n-1}(n-1)!}{(2n-1)!}\{(\tfrac{1}{2}a+\tfrac{1}{2})_n + (\tfrac{1}{2}a)_n\}, \tag{3.23} $$

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n-1 \end{array};2\right] = \frac{2^{2n+1}n!}{(2n+1)!}(\tfrac{1}{2}a+\tfrac{1}{2})_{n+1}, \tag{3.24} $$

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n+2 \end{array};2\right] = \frac{2^{2n-1}(n-2)!}{(2n-2)!}\left\{\frac{1-a-n}{1-a-2n}(\tfrac{1}{2}a+\tfrac{1}{2})_n + (\tfrac{1}{2}a)_n\right\}, \tag{3.25} $$

$$ {}_2F_1\left[\begin{array}{c} -n, a \\ -2n-2 \end{array};2\right] = \frac{2^{2n+3}n!}{(2n+2)!}\left\{\frac{(1-a-n-j)}{1-a-2n-2j}(\tfrac{1}{2}a+\tfrac{1}{2})_{n+2} - (\tfrac{1}{2}a)_{n+2}\right\} \tag{3.26} $$

and so on.

The above evaluations agree with those given in [2, 3], although presented in a different format; the results (3.1) and (3.6), together with (3.11), (3.12) and (3.17), are also recorded in [**?**] in another form.

4. An application of Theorem 1

Kummer's second theorem applied to the confluent hypergeometric function ${}_1F_1$ is [10, p. 12]

$$ e^{-x/2}{}_1F_1\left[\begin{array}{c} a \\ 2a \end{array};x\right] = {}_0F_1\left[\begin{array}{c} - \\ a+\frac{1}{2} \end{array};\frac{x^2}{16}\right] = (\tfrac{1}{4}x)^{\frac{1}{2}-a}\Gamma(a+\tfrac{1}{2})I_{a-\frac{1}{2}}(\tfrac{1}{2}x), \tag{4.1} $$

where $I_\nu(z)$ denotes modified Bessel function of the first kind. We now show how the result in Theorem 1 can be used to derive a generalization of the above theorem for the functions

$$ e^{-x/2}{}_1F_1\left[\begin{array}{c} a \\ 2a \pm j \end{array};x\right] $$

for arbitrary integer j.

We have upon series expansion

$$ e^{-x/2}{}_1F_1\left[\begin{array}{c} a \\ 2a \pm j \end{array};x\right] = \sum_{n=0}^{\infty}\frac{(-\frac{1}{2}x)^n}{n!}\sum_{m=0}^{\infty}\frac{(a)_m}{(2a\pm j)_m}\frac{x^m}{m!} = \sum_{n=0}^{\infty}\sum_{m=0}^{\infty}\frac{(-1)^n(a)_m x^{m+n}}{2^n(2a\pm j)_m m!n!}. $$

Making the change of summation index $n \to n-m$ and using the fact that $(n-m)! = (-1)^m m!/(-n)_m$, we find

$$ e^{-x/2}{}_1F_1\left[\begin{array}{c} a \\ 2a \pm j \end{array};x\right] = \sum_{n=0}^{\infty}\sum_{m=0}^{n}\frac{(-1)^n(a)_m(-n)_m x^n}{2^{n-m}(2a\pm j)_m m!n!} $$

$$ = \sum_{n=0}^{\infty}\frac{(-\frac{1}{2}x)^n}{n!}\,{}_2F_1\left[\begin{array}{c} -n, a \\ 2a \pm j \end{array};2\right]. $$

Separation of the above sum into even and odd n, use of the evaluation of the ${}_2F_1(2)$ series given in (2.1) and (2.2) followed by inversion of the order of summation then leads to the result

$$ e^{-x/2}{}_1F_1\left[\begin{array}{c} a \\ 2a \pm j \end{array};x\right] = \sum_{r=0}^{j_0}(-1)^r\binom{j}{2r}\sum_{n=0}^{\infty}\frac{(-n)_r(a+\delta_j)_{n-r}x^{2n}}{2^{2n}(2a\pm j)_{2n}n!} $$

$$\mp \sum_{r=0}^{j_0} (-1)^r \binom{j}{2r+1} \sum_{n=0}^{\infty} \frac{(-n)_r (a+\delta_j)_{n-r} x^{2n+1}}{2^{2n+1} (2a \pm j)_{2n+1} n!}, \qquad (4.2)$$

where j_0 and δ_j are defined in Theorem 1. When $j = 0$ it is easily seen that (4.2) reduces to (4.1).

The result (4.2) is given in a different form in terms of modified Bessel functions in [7, Vol. 3, p. 579].

Acknowledgement: One of the authors (YSK) acknowledges the support of the Wonkwang University Research Fund (2014).

References

[1] J. Choi, A. K. Rathie and H. M. Srivastava, A generalization of a formula due to Kummer, Integral Transforms and Special Functions, **22**(11) (2011) 851–859.

[2] W. Chu, Terminating hypergeometric ${}_2F_1(2)$ series, Integral Transforms and Special Functions **22**(2) (2011) 91–96.

[3] Y. S. Kim and A. K. Rathie, Some results for terminating ${}_2F_1(2)$ series, J. Inequal. Applications, 2013:365 (2013).

[4] Y. S. Kim, M. A. Rakha and A. K. Rathie, Generalization of Kummer's second theorem with applications, Comput. Math. Math. Phys. **55**(3) (2010) 387–402.

[5] Y. S. Kim, A. K. Rathie and R. B. Paris, Generalization of two theorems due to Ramanujan, Integral Transforms and Special Functions **24**(4) (2013) 314–323.

[6] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clark (eds.), NIST Handbook of Mathematical Functions, Cambridge University Press, Cambridge, 2010.

[7] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, Integrals and Series: Special Functions, Gordon and Breach, New York, 1988.

[8] M. A. Rakha and A. K. Rathie, Generalizations of classical summation theorems for the series ${}_2F_1$ and ${}_3F_2$ with applications, Integral Transforms and Special Functions **22**(11) (2011) 823–840.

[9] A. A. Samoletov, A sum containing factorials, J. Comput. Appl. Math. **131** (2001) 503–504.

[10] L. J. Slater, Confluent Hypergeometric Functions, Cambridge University Press, Cambridge, 1960.

[11] L. J. Slater, Generalized Hypergeometric Functions, Cambridge University Press, Cambridge, 1966.

[12] H. M. Srivastava, Remarks on a sum containing factorials, J. Comput. Appl. Math. **142** (2002) 441-444.